



Original Article

Effects of Local Transplantation of Autologous Bone Marrow Mesenchymal Stem Cells in Combination with Low Level Laser Therapy in Repair of Experimental Acute Spinal Cord Injury in Rats

Seyed Sadra Izadi¹, Mohammad Mehdi Dehghan^{1,2*}, Davood Sharifi¹, Seyed Mehdi Nassiri³, Seyed Hossein Marjanmehr⁴, Maryam Zabihi¹, Javad Ashrafi-Helan⁵, Hessam Akbarein⁶

Abstract

Objective- The aim of this study was to demonstrate the efficacy of MSCs transplantation in combination with low level laser irradiation (LLLI) in repair of experimental acute spinal cord injury.

Design- Experimental study.

Animals- 28 adult male Wistar Rats.

Procedures- A balloon- compression technique was used to produce an injury at the T8-T9 level of spinal cord applying Fogarty embolectomy catheter. In group-1, the autologous MSCs were transplanted to the spinal cord lesion and followed by treatment with low level laser irradiation during 15 consecutive days in group-2. The injured rats in third group were treated by LLLI alone. The functional recovery was assessed using the Basso-Beattie-Bresnahan (BBB) locomotion scoring within 5 weeks.

Results- In these three treatment groups, the scores were significantly higher than control group. The differences between group-2 and two other treatment groups were statistically significant during all five weeks after treatment. There were no significant differences in BBB score between group-1(MSCs) and group-3(LLLI) at 3rd, 4th and 5th weeks of treatment. According to the histopathological findings, the best response was observed in group-2 (MSCs+LLLI) that repair of injured parts of dorsal funiculi and less cavitation were occurred by proliferation of mesenchymal stem cells and their differentiation to glial cells especially oligodendrocytes resulting in axon regeneration and relatively spinal cord recovery.

Conclusion and Clinical Relevance- The findings of present study demonstrated that concurrent use of LLLI and local transplantation of MSCs exhibited profound effects on axon regeneration and revealed remarkable functional improvement. These results suggested that MSCs characteristics could be influenced by low level laser irradiation, so this treatment might be as a useful procedure for neural regeneration, although further detailed investigations need to be carried out particularly in clinical cases.

Keywords- Spinal cord injury, Mesenchymal stem cells, Low level laser therapy

Received: 12 May 2016; **Accepted:** 11 September 2017; **Online:** 31 October 2017

Introduction

Spinal cord injury (SCI) is a trauma-induced disease state, happening in the central nerves system (CNS). SCIs cause variant degrees of decimation and

functional loss to the axons of the spine and their downstream targets.¹

The majority of SCIs are due to preventable causes including road traffic crashes, violence or falls.² The CNS neurons have the characteristic ability to regenerate a lost axon. Growth inhibitory molecules, lack of appropriate trophic support, and immune system reactions are ascribed factors to axon regeneration failure after SCI.³ Spinal cord injuries are accompanied by a number of complications, causing death of neurons, degeneration of nerve fibers, hemorrhage, and eventually the absence of complete regeneration in areas of injury. In most of the cases, traditional methods of treatment are very rarely able to restore the lost functions of tissues. However, the use of stem cells in these patients gives hope for the

¹Department of Veterinary Surgery & Radiology, Faculty of Veterinary Medicine, University of Tehran, Tehran- Iran.

²Institute of Biomedical Research, University of Tehran, Tehran, Iran.

³Department of Internal Medicine, Faculty of Veterinary Medicine, University of Tehran, Tehran- Iran.

⁴Department of Pathology, Faculty of Veterinary Medicine, University of Tehran, Tehran- Iran.

⁵Department of Pathobiology, Faculty of Veterinary Medicine, University of Tabriz, Tabriz- Iran.

⁶Department of Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, University of Tehran, Tehran- Iran.

Address all correspondence to Mohammad Mehdi Dehghan (DVM, DVSc) E-mail:mdehghan@ut.ac.ir

www.ivsajournals.com © Iranian Journal of Veterinary Surgery, 2017

This work is licensed under the terms of the Creative Commons Attribution (CC BY-NC-SA 4.0)

DOI: 10.22034/ivsa.2017.51719

opportunity to achieve functional improvements.⁴ To date there is insufficient evidence that would support approved clinical treatments to form neurons and axons immediate protection or to increase their regeneration, but using stem cell therapy in patients with SCI gives hope for opportunity to achieve functional improvement and constitutes a target of great unmet medical demand.⁵ Among many treatments being developed for SCI, the placement of different types of grafts embedded with cell populations into areas of damage has been one of the most commonly regenerative approaches, attracting scientists and physicians for the last 15 years.⁶ Many kinds of somatic cells have been studied as transplants for the treatment of SCI, which include bone marrow stromal cells (BMSCs)⁷, olfactory ensheathing cells⁸, Schwann cells⁷, dental pulp-derived cells⁹, epidermal neural crest stem cells¹⁰, skin-derived precursor cells¹¹, adipose-derived stromal cells¹², and choroid plexus epithelial cells.¹³ MSCs have been being applied in treatment of different disorders including spinal cord injury in animal models.¹⁴ Oliveri *et al.* in a systematic review with meta-analyses suggested that recovery of locomotor function in traumatic SCI animal models can be promoted by MSCs therapy.¹⁵ Also, it is assumed that implantation of MSCs will have better effect than injection them into blood or cerebrospinal fluid, since it will increase amount of MSCs in damaged site. On the other hand, the use of electrotherapeutics modalities is a common practice in physiotherapy with regenerative purposes, especially using low-level laser irradiation (LLLI) holds promise for future of tissue engineering and regenerative medicine.¹⁵⁻¹⁷ LLLI can cause in prevention of MSCs apoptosis and improvement of MSCs proliferation, migration, and adhesion at low-powers and low-levels of red/close infrared light enlightenment¹⁸, which is approved as a dose dependant procedure.¹⁹ Hou *et al.* documented that LLLI could increase growth factors secretion, stimulate proliferation and facilitate myogenic differentiation of BMSCs. Accordingly; LLLI may give a novel way to deal with the preconditioning of BMSCs in vitro before transplantation.²⁰

Thus, this study was conducted to evaluate the effect of transplantation of BMSCs associated in combination with LLLI for the treatment of the contusion-injured spinal cord of rats.

Material and Methods

The study was conducted with 28 male Fischer-344 Wistar rats, 8 to 12 weeks of age. In order to decrease the variability of spinal canal size, only animals with body weights between 300-350 g were included. All animals were kept in large and well-lit plastic containers. These containers were kept separately and

at laboratory controlled temperature of 21 °C. Additionally, the containers were maintained with a daily photoperiod of 12 hour of light for seven days. Each animal had free access to food and water *ad libitum*. After 7-day adaptation period, bone marrow was extracted from femur bones of rats. MSCs isolation and propagation lasted a total of 3 weeks. At this time, spinal cord injury was induced in rats. One week after induction of SCI, injection of undifferentiated autologous MSCs was performed by using a Hamilton Syringe. One day after MSCs transplantation, laser therapy was started by a low level laser with a wavelength of 780 nm and a power of 250 mw. Two weeks later, Basso- Beattie-Bresnahan (BBB) functional scoring test was used for assessing the locomotor capacity of rats after SCI, and continued weekly for six weeks. Finally, histopathological evaluations were performed on the histopathological samples of the injured region (Fig 1).²¹




Figure 1 Schematic drawing showing the induction of spinal cord injury (SCI); and local transplantation of MSCs in group-1, local transplantation of MSCs followed by low level laser irradiation (LLLI) in group-2; LLLI in group-3 and Control group as group-4 of experimental animals.

Isolation of bone marrow stromal cells (BMSCs)

MSCs were harvested from the femoral bone marrows of rats and then the cells were transplanted into the same rat in order to decrease the chance of rejection. Procedures of extraction, isolation, and propagation of BMSCs were as same as Wakitan.²² In summary, MSCs were harvested by FNA (fine needle aspiration) technique from the femoral bone marrow. Rats were anesthetized by intramuscular injection of Ketamine HCl and Xylazine at 75 mg/kg and 10 mg/kg, respectively. Following a 5mm length skin incision a small opening (1-1.5 mm) in the femur was drilled. Then, a 2 ml syringe with a 21 G needle containing 500-750 IU heparin was used to aspirate of 0.5-1 ml of bone marrow. The samples were diluted with L-15 medium (2 ml) containing 3 ml of Ficoll. Then, samples

were centrifuged (2,000 rpm) for 15 minutes, then cells in the mononuclear layer were harvested and were re-suspended in 2 ml serum-free medium, then centrifuged (2,000 rpm) for 15 minutes and were re-suspended in 1 ml DMEM.

Spinal cord injury

Induction of SCI in rats was performed by a method described by Vanicky et al.²³ In brief, rats were anesthetized as described above. When they were in stable situation, a midline incision of 2 cm was created over the T10-L1 spinous processes, under sterile conditions. Then the spinous processes of vertebrae T10 – T11 were removed following the dissection of the regional skin and muscles. Under magnification, vertebral arch of T10 was drilled using a micromotor bur. A groove was also drilled in the midline on the dorsal surface of T11 vertebral lamina to guide the insertion of the catheter and keep it positioned in the midline. A saline loaded 2-French Fogarty catheter was linked to an airtight 50- μ l Hamilton syringe held in a precise sampling device. After insertion of the catheter into epidural space in a way that the center of the balloon was rested at T8-T9 level of spinal cord, the balloon was distended quickly with 20 μ l volume of saline for 5 minutes. Then, serum was removed from the catheter and catheter was pulled out slowly. Skin and the other layers were attached together by appropriate suture placement in anatomical layers. Bladder was evacuated manually at least twice a day until reflex bladder was approved. Antibiotic therapy was performed by Enrofloxacin administration (10 mg/kg, every 24 hour) for one week. All rats were paraplegic after injury, with no signs of functional recovery. All experiments were carried out in accordance with approved guidelines of the Iran Animal Care Committee and were approved by the Faculty of Veterinary Medicine, University of Tehran Animal Care Committee.

MSCs transplantation

Seven days after induction of SCI, rats were anesthetized again as described before and vertebral arches of the T8-T9 were removed using micro-motor and a burr (Stryker Corporation, USA). Injured rats were treated with 1×10^6 autologous undifferentiated BMSCs by insertion the tip of a 20- μ l Hamilton syringe through the intact dura. Tip of the syringe were inserted into the center of the developing lesion cavity 3 mm

cranial and 3mm caudal to the cavity (penetration depth of 1.0 mm at an angle of 40–45° past perpendicular).

Low- Level Laser Irradiation

One day after MSCs transplantation, LLLI procedure has performed in group2 (MSCs+Laser) and group3 (Laser without MSCs) as described below. Briefly, groups 2 and 3 of paralyzed rats were irradiated with red or near-infrared laser via transcutaneous application.²⁴ LLLI was started immediately one day after surgery and was continued daily for two weeks. A 250mW NIR laser (wavelength 780; continuous wave (CW)) was transcutaneously irradiated over the 1cm distance between T8 and T10 for 15 min daily (spot size 3 mm², laser fluence 10 J/cm²).²⁵

Behavioral Assessment

During the 6-week follow up, motor activity of hind limbs was evaluated according to Basso-Beattie-Bresnahan (BBB) open field locomotor rating scale.²⁶ BBB scores include a wide range from 0-21 (0: no observable movement in hind limbs; 21: normal locomotion). These scales represent the recovery of locomotor activity and are categorized according to the sum of the animal's joint-movements, hind limb movements, stepping, forelimb and hind limb coordination, trunk position and stability, paw placement and tail position.²⁷ All of the rats movements were recorded for better analysis and more detailed assessment. Sensory recovery was evaluated by pinch test and observation of withdrawal reflex. This test gives useful information about distance and rate of regeneration of sensory neurons. In summary, rats were placed on examination desk in a way that the muscles of the hind limbs were completely relaxed (knee flexed at 130° and ankle at 90°) using anatomic supports.²⁸ Then, to evaluate nociceptive withdrawal reflex, Kelly forceps were placed on the rats hind limb fingers and gradual pressure was applied until the animal showed any aversive response such as withdrawal of the limb, vocalization, and struggling.²⁹

Histopathological assessment

Five weeks after implantation of the undifferentiated BMSCs in spinal cord, the rats were deeply anesthetized by injection of 100 mg/kg pentobarbital sodium intraperitoneally, and after that perfused transcardially with 200 ml 0.1 M phosphate buffer (pH 7.4) continued by 300 ml 4% phosphate-buffered saline (pH 7.4) containing 4% paraformaldehyde and 1% glutaraldehyde. Spinal cords were sectioned transversely

from T7 to T10.²¹ The injured region of the spinal cord was fixed in formalin 10% and then tissue sections were obtained from it. After fixation, transverse sections of spinal cord at T7 to T10 were embedded, cut into 5 µm thick sections, and stained using hematoxylin and eosin. Afterwards, histopathological assessment of cells, myelinated, and demyelinated neural fibers were performed under 40, 100, 200 magnifications by a pathologist blind to the groups.²¹

It is certified that all the animal experiments followed the applicable institutional and governmental regulations concerning the ethical use of animals.

Statistical Analysis

Statistical analyses were done by SPSS package Version 19.0. The data were described by Mean±SEM. The data were analyzed by One-Way ANOVA and Tukey Post Hoc tests. A *p*-value less than 0.05 were statistically considered significant.

Results

Behavioral analysis

During the 6-week follow up, the locomotor and sensory recovery of rats was weekly evaluated by two observers. Rats in all groups showed no significant locomotion in hind limbs one week after induction of SCI. Three weeks after induction of SCI (two weeks after beginning of therapies), BBB score of group-1 (MSCs) was increased to 4.88. At the same time, in the group-2 (MSCs + LLLI) and group-3 (LLLI), BBB scores were increased to 10.81 and 6.38, respectively. Six weeks after induction of SCI (after five weeks of therapy and/or at the end of behavioral assessment), BBB score of group-1 (MSCs) increased to 8.25. At the same time, BBB score increased to 17.13 and 8.31 for the group-2 (MSCs + LLLI) and group-3 (LLLI) respectively. At week-6, the differences among group-2 (MSCs + LLLI) and two other groups (MSCs and LLLI) were statistically significant (*p*<0.05). There were no significant differences in BBB score between group-1 (MSCs) and group-3 (LLLI) at week-3, week-4 and week-5 of therapy period (Fig 2).

Histopathological findings

In the control group, the dorsal funiculi of spinal cord were edematous with focal destruction and degeneration of myelin, swelling of axons and microcavitations was apparently seen. Proliferation




Figure 2. Mean±SEM of Basso-Beattie-Bresnahan (BBB) test results within six weeks after SCI

of astrocytes, microglial cells and especially oligodendrocytes were present. Some of microglial cells and large foam cells were seen around of destructed areas. Gray matter especially in dorsal horns of spinal cord was atrophic with a severe hypo-cellularity of neurons. The number of these large neurons was greatly decreased from 200 neurons in intact part of spinal cord to 30 neurons in injured areas.

According to histopathological evaluation, the best response was observed in the group that treated with combination of MSCs and LLLI. In this group, repair of injured parts of dorsal funiculi was occurred by proliferation of mesenchymal cells and differentiation of them to glial cells especially oligodendrocytes. It caused promotion of axon regeneration and relatively spinal cord recovery but hypocellularity of dorsal horn was apparent. There were no acute inflammatory reaction and granulomatous reaction. In laser and MSCs groups, evidences of focal destruction of dorsal funiculi and foam cells, astrocytosis and astrogliosis were still seen. In these groups, repair of injured parts of dorsal funiculi was not completed and proliferation of mesenchymal cells and differentiation of them to glial cells especially oligodendrocytes were mild to moderate. Axons degeneration and hypocellularity of dorsal horn was apparent. There were no acute inflammatory reactions and granulomatous reactions (Fig 3).

Discussion

Results of the BBB locomotors scoring test indicated that concurrent use of laser irradiation associated with undifferentiated BM-MSCs in spinal cord injured rats has better impact than use of laser therapy or BM-MSCs alone. Also it was obvious and believed that use of laser therapy or BMSCs could improve cellular structure of spinal cord and finally locomotor and sensory recovery. Although, MSCs therapy and LLLI could promote axons regeneration and recovery of injured spinal cord,




Figure 3. Histopathological findings in different groups: A, Control Group; B, BMSCs Group; C, LLLI Group; D, BMSCs + LLLI Group. **A-1)** A microscopic view of spinal cord in control group, dorsal funicular is edematous with a focal destruction (a), degeneration of myelin and microcavitations (b) is apparently seen especially around of destructed area. Gray matter of dorsal horn shows severe hypo-cellularity (c) of neurons (H&E, $\times 100$). **A-2)** A microscopic view of spinal cord in control group, dorsal funicular is edematous with a focal destruction (a), degeneration of myelin (a), swelling of axons (d) and microcavitations (b) is apparently seen especially around of destructed area (H&E, $\times 200$). **B-1)** A microscopic view of spinal cord in BMSCs group, in these groups, repair of injured parts of dorsal funicular is not completed and proliferation of mesenchymal cells and differentiation of them to glial cells (e) is mild to moderate. Axons degeneration (d) and hypocellularity (c) of dorsal horn is apparent (H&E, $\times 100$). **B-2)** A closer microscopic section of spinal cord in BMSCs group, repair of injured parts of dorsal funicular is relatively completed and proliferation of mesenchymal cells and differentiation of them to glial cells, axon regeneration (f) and spinal cord recovery is apparent (H&E, $\times 200$). **C-1)** A microscopic view of spinal cord in LLLI group, in these groups, repair of injured parts of dorsal funicular is not completed. Axons degeneration (d) and hypocellularity (c) of dorsal horn is apparent (H&E, $\times 100$). **C-2)** A closer microscopic section of spinal cord in LLLI group, repair of injured parts of dorsal funicular is relatively completed and proliferation of glial cells (e) and axon regeneration (f) is apparent (H&E, $\times 200$). **D-1)** A microscopic section of spinal cord in BMSCs + LLLI group, repair of injured parts of dorsal funicular is completed and axon regeneration (f) and spinal cord recovery is apparent but dorsal horn is hypocellular (c) (H&E, $\times 100$). **D-2)** A closer microscopic view of spinal cord in BMSCs + LLLI group, repair of injured parts of dorsal funicular is completed and axon regeneration (f) and spinal cord recovery is apparent. In this group, repair of injured parts of dorsal funicular is occurred by proliferation of mesenchymal cells and differentiation of them to glial cells (H&E, $\times 200$).

as an independent treatment method, but concurrent use of low level laser therapy and local transplantation of BM-MSCs revealed remarkable improvement of locomotion recovery in the rats with spinal cord injury. Also it was shown that LLLI played a major role to achieve better functional improvement and accelerated functional recovery specially at the end of the week 2 in combination of local transplantation of MSCs concurrently.

Among several experimental strategies which have been investigated for spinal cord injuries treatment, cell therapy seems that has the best results for improving clinical situation of a paralytic patient.³⁰ BMSCs appear to be one of the best candidates among various types of cells utilizing for this purpose.³¹ Application of BMSc as a treatment of many diseases such as osteogenesis imperfecta, mucopolysaccharidoses, graft versus host disease, and myocardial infarction has been evaluated and the results were remarkable.³² Masayoshi Ohta et al. (2004), showed that BMSCs can exert direct influence by reduction in volume of injured-spinal cord cavities during first three weeks after cell transplantation. In addition, transplantation of BMSCs exhibit profound effects by releasing some beneficial substances into the CSF, resulting in the repair of the spinal cord lesions.³³ These findings suggest that production and secretion of some trophic factors and their synchronized actions are beneficial for neurons tissue repair; such as Colony Stimulating Factor-1 (CSF-1), interleukins, stem cell factors^{34,35}, BDNF, NGF, HGF and VEGF.³⁶ It also was reported that glial cells which are stimulated by BMSCs produce some neurotrophic substances such as BDNF and NGF^{35,37,38}, resulting in repair of the lesions and functional improvement, subsequently.³³ Recently MSCs have shown different properties, for examples, anti-apoptotic, immunomodulatory, anti-inflammatory, angiogenic and trophic impacts. These capacities are accepted to be interceded by transient paracrine by-stander mechanisms and cell-to-cell contact in response to the local damaged host tissue environment rather than cell replacement and long-term cell engraftment.¹⁵ Among different kinds of MSCs, BMSCs are remained the 'gold standard' having been most described in preclinical and clinical studies. BMSCs can be simply isolated and expanded in vitro to several hundred millions cells within a generally brief timeframe.¹⁵ On the other hand, the results shown that laser therapy can significantly improve locomotor function of hind limbs.

It has been demonstrated that LLLI promotes differentiation and proliferation of human osteoblast cells, in vitro³⁹, proliferation of cardiac and mesenchymal stem cells in culture at 1 J/cm² and 3 J/cm².⁴⁰ Studies on LLLI have indicated that it could play a major role in many tissue regenerating processes such as wound healing^{41,42}, fibroblast proliferation⁴³, nerve regeneration⁴⁴, and collagen synthesis and could increase migration of stem cells in vitro by changing the metabolism of stem cells and increasing the adenosine triphosphate level of MSCs.²⁰ Furthermore, LLLI can improve proliferation of rat mesenchymal bone marrow and cardiac stem cells in vitro.⁴⁰ It is remarkable that laser irradiation at 665-675 nm wave length induces the maximum effect on cell proliferation and growth factors secretion in vitro, whereas irradiation at 810 nm (or higher) inhibited cell division in vitro.⁴⁵ LLLI can promote cell proliferation and increase, DNA, and RNA synthesis and collagen synthesis which can initiate mitosis in cultured cells. Also, mitochondrial and cell membrane receptors can be stimulated by LLLI. These receptors can convert the light energy into chemical energy (ATP) which increases cell proliferation rate and improves cellular functions.⁴⁵ Finally recent studies demonstrated that transcutaneous application of a 780 nm laser can improve the axonal regeneration and functional recovery.^{46,47} Therefore, it is assumed laser therapy can affect the grafted BMSCs and amplify their function. Based on histopathological evaluation, in this study, the best response was observed in the group treated with combination of laser-cell therapy. In this group, repair of injured parts of dorsal funiculi was occurred by proliferation of mesenchymal cells and differentiation of them to glial cells, especially oligodendrocytes. It resulted in promotion of axon regeneration and relatively spinal cord recovery.

Acknowledgments

This research was supported by grant from University of Tehran. The authors are grateful to Dr M S Pedram, Dr M S Ghodrati, Dr H Reisdanaei, Mr Isanejad and Mr Baninajar for their technical helps and advices.

Conflicts of interest

None

References

1. Finnegan J and YE H. Cell therapy for spinal cord injury informed by electromagnetic waves. *Regenerative Medicine*, 2016;11(7):675-691.
2. World Health Organization Web site. Spinal cord

- injury. *fact sheet* (2013). available at: <http://www.who.int/mediacentre/factsheets/fs384/en/>
3. Hofstetter CP, Schwarz EJ, Hess D, Widenfalk J, El Manira A, Prockop DJ and Olson L. Marrow stromal cells form guiding strands in the injured spinal cord and promote recovery. *Proceedings of the National Academy of Sciences*, 2002;99(4):2199-2204.
 4. Kakabadze Z, Kipshidze N, Mardaleishvili K, Chutkerashvili G, Chelishvili I, Harders A, Loladze G, Shatirishvili G, Kipshidze N, Chakhunashvili D, Chutkerashvili K. Phase 1 trial of autologous bone marrow stem cell transplantation in patients with spinal cord injury. *Stem Cells International*, 2016;6768274. doi: 10.1155/2016/6768274.
 5. Lu p, Ahmad R, and Tuszynski MH. Neural Stem Cells for Spinal Cord Injury. In Tuszynski MH, ed. *Translational Neuroscience Fundamental Approaches for Neurological Disorders*. Springer, 2016;297-314.
 6. Jendelova P, Machova-Urdzikova L and Sykova E. The role of mesenchymal stromal cells in spinal cord injury. In. Atkinson K, ed. *The biology and therapeutic application of mesenchymal cells*. WILEY Blackwell, 2017;714-729.
 7. Ide C, Nakano N and Kanekiyo K. Cell transplantation for the treatment of spinal cord injury—bone marrow stromal cells and choroid plexus epithelial cells. *Neural Regeneration Research*, 2016;11(9): 1385-1388.
 8. Tharion G, Indirani K, Durai M, Meenakshi M, Devasahayam SR, Prabhav NR, Solomons C, Bhattacharji S. Motor recovery following olfactory ensheathing cell transplantation in rats with spinal cord injury. *Neurology India*, 2011;59(4):566.
 9. Sakai K, Yamamoto A, Matsubara K, Nakamura S, Naruse M, Yamagata M, Sakamoto K, Tauchi R, Wakao N, Imagama S, Hibi H, Kadomatsu K, Ishiguro N, Ueda M. Human dental pulp-derived stem cells promote locomotor recovery after complete transection of the rat spinal cord by multiple neuro-regenerative mechanisms. *The Journal of clinical investigation*, 2012;122(1):80-90.
 10. Gericota B, Anderson J S, Mitchell G, Borjesson D L, Sturges B K, Nolte J A, and Sieber-Blum M. Canine epidermal neural crest stem cells: characterization and potential as therapy candidate for a large animal model of spinal cord injury. *Stem cells translational medicine*, 2014;3(3):334-345.
 11. Biernaskie J, Sparling J S, Liu J, Shannon C P, Plemel J R, Xie Y and et al. Skin-derived precursors generate myelinating Schwann cells that promote remyelination and functional recovery after contusion spinal cord injury. *Journal of Neuroscience*, 2007;27(36):9545-9559.
 12. Arboleda D, Forostyak S, Jendelova P, Marekova D, Amemori T, Pivonkova H, Masinova K, Sykova E. Transplantation of predifferentiated adipose-derived stromal cells for the treatment of spinal cord injury. *Cellular and Molecular Neurobiology*, 2011;31(7):1113-1122.
 13. Kanekiyo K, Nakano N, Noda T, Yamada Y and Suzuki Y. Transplantation of choroid plexus epithelial cells into contusion-injured spinal cord of rats. *Restorative neurology and neuroscience*, 2016;34(3):347-366.
 14. Sobhani A, Khanlarkhani N, Baazm M, Mohammadzadeh F, Najafi A, Mehdinejadi F S A. Multipotent Stem Cell and Current Application. *Acta Medica Iranica*, 2017, 55.1: 6-23 Multipotent Stem Cell and Current Application, 2017;55: 6-23.
 15. Oliveri R S, Bello S and Biering-sørensen F. Mesenchymal stem cells improve locomotor recovery in traumatic spinal cord injury: systematic review with meta-analyses of rat models. *Neurobiology of disease*, 2014;62:338-353.
 16. Paula A A, Nicolau R A, de Oliveira Lima M, Salgado M A C, and Cogo J C. Low-intensity laser therapy effect on the recovery of traumatic spinal cord injury. *Lasers in medical science*, 2014;29(6):1849-1859.
 17. Hou J F, Zhang H, Yuan X, Li J, Wei Y J and H S S. In vitro effects of low-level laser irradiation for bone marrow mesenchymal stem cells: Proliferation, growth factors secretion and myogenic differentiation. *Lasers in surgery and medicine*, 2008;40(10):726-733.
 18. Huang Y Y, Chen A C H, Carroll J D and Hamblin M R. Biphasic dose response in low level light therapy. *Dose-Response*, 2009;7(4):358-383.
 19. Bolton P, Young S and Dyson M. The direct effect of 860 nm light on cell proliferation and on succinic Dehydrogenase Activity Of Human Fibroblasts In vitro. *Laser Therapy*, 1995;7(2):55-60.
 20. Hou J, Zhang H, Yuan X and Li J. In vitro effects of low-level laser irradiation for bone marrow mesenchymal stem cells: Proliferation, growth factors secretion and myogenic differentiation. *Lasers in surgery and medicine*, 2008;40(10):726-733.
 21. Pedram M S, Dehghan M M, Soleimani M, Sharifi D, Marjanmehr S H and Nasiri Z. Transplantation of a combination of autologous neural differentiated and undifferentiated mesenchymal stem cells into injured spinal cord of rats. *Spinal Cord*, 2010;48(6):457-463.
 22. Wakitani S, Saito T and Caplan A. I. Myogenic cells derived from rat bone marrow mesenchymal stem cells exposed to 5-azacytidine. *Muscle & nerve*, 1995;18(12):1417-1426.
 23. Vanický I, Urdzíkova, L, Saganová K, Čížková D and Gálík J. A Simple and Reproducible Model of

Spinal Cord Injury Induced by Epidural Balloon Inflation in the Rat. *Journal of Neurotrauma*, 2001;18(12):1399-407.

24.Nissan M, Rochkind S, Razon N and Bartal A. HeNe laser irradiation delivered transcutaneously: its effect on the sciatic nerve of rats. *Lasers in surgery and medicine*, 1986;6(5):435-438.

25.Rochkind S, Shahar A, Alon M and Nevo Z. Transplantation of embryonal spinal cord nerve cells cultured on biodegradable microcarriers followed by low power laser irradiation for the treatment of traumatic paraplegia in rats. *Neurological research*, 2002;24(4):355-360.

26.Basso D M, Beattie M S and Bresnahan J C. A sensitive and reliable locomotor rating scale for open field testing in rats. *Journal of neurotrauma*, 1995;12(1)1-21.

27.Segal Y, Segal L, Shohami E, Sasson E, and Blumenfeld-Katzir T, Cohen A, Levy A and Alter A. The Effect of Electromagnetic Field Treatment on Recovery from Spinal Cord Injury in a Rat Model—Clinical and Imaging Findings. *International Journal of Neurorehabilitation*, 2016;3: 203. doi:10.4172/2376-0281.1000203.

28.Perrotta A, Chiacchiaretta P, Anastasio M G, Pavone L, Grillea G, Bartolo M, Siravo I E, Colonnese C, De Icco R, Serrao M, Sandrini G, Pierelli F, Ferretti A. Temporal summation of the nociceptive withdrawal reflex involves deactivation of posterior cingulate cortex. *European Journal of Pain*, 2017;21(2):289-301.

29.Hayes RL, Bennett GJ, Newlon PG and Mayer DJ. Behavioral and physiological studies of non-narcotic analgesia in the rat elicited by certain environmental stimuli. *Brain research*, 1978;155(1):69-90.

30.McKerracher L. Spinal cord repair: strategies to promote axon regeneration. *Neurobiology of disease*, 2001;8(1):11-18.

31.Deng W, Obrocka M, Fischer I and Prockop D J. In vitro differentiation of human marrow stromal cells into early progenitors of neural cells by conditions that increase intracellular cyclic AMP. *Biochemical and biophysical research communications*, 2001;282(1):148-152.

32.Prockop DJ and Olson S D. Clinical trials with adult stem/progenitor cells for tissue repair: let's not overlook some essential precautions. *Blood*, 2007;109(8):3147-3151.

33.Ohta M, Suzuki Y, Noda T, Ejiri Y, Dezawa M, Kataoka K and et al. Bone marrow stromal cells infused into the cerebrospinal fluid promote functional recovery of the injured rat spinal cord with reduced cavity formation. *Experimental*

neurology, 2004;187(2):266-278.

34.Eaves CJ, Cashman JD, Kay RJ, Dougherty GJ, Otsuka T, Gaboury LA, Hogge DE, Lansdorp PM, Eaves AC, Humphries RK. Mechanisms that regulate the cell cycle status of very primitive hematopoietic cells in long-term human marrow cultures. II. Analysis of positive and negative regulators produced by stromal cells within the adherent layer. *Blood*, 1991;78(1):110-117.

35.Majumdar MK, Thiede MA, Mosca JD, Moorman M and Gerson SL. Phenotypic and functional comparison of cultures of marrow-derived mesenchymal stem cells (MSCs) and stromal cells. *Journal of cellular physiology*, 1998;176(1):57-66.

36.Chen X, Katakowski M, Li Y, Lu D, Wang L, Zhang L, Chen J, Xu Y, Gautam S, Mahmood A, Chopp M. Human bone marrow stromal cell cultures conditioned by traumatic brain tissue extracts: Growth factor production. *Journal of neuroscience research*, 2002;69(5):687-691.

37.Mahmood A, Lu D, Wang L and Chopp M. Intracerebral transplantation of marrow stromal cells cultured with neurotrophic factors promotes functional recovery in adult rats subjected to traumatic brain injury. *Journal of neurotrauma*, 2002;19(12):1609-1617.

38.Wang L, Li Y, Chen J, Gautam SC, Zhang Z, Lu M, and Chopp M. Ischemic cerebral tissue and MCP-1 enhance rat bone marrow stromal cell migration in interface culture. *Experimental hematology*, 2002;30(7):831-836.

39.Stein A, Benayahu D, Maltz L and Oron U. Low-level laser irradiation promotes proliferation and differentiation of human osteoblasts in vitro. *Photomedicine and Laser Therapy*, 2005;23(2):161-166.

40.Tuby H, Maltz L and Oron U. Low-level laser irradiation (LLLI) promotes proliferation of mesenchymal and cardiac stem cells in culture. *Lasers in surgery and medicine*, 2007; 39(4):373-378.

41.Beckmann K H, Meyer-Hamme G, Schröder S and der S. Low level laser therapy for the treatment of diabetic foot ulcers: a critical survey. *Evidence-Based Complementary and Alternative Medicine*, vol. 2014, Article ID 626127, 9 pages, 2014. doi:10.1155/2014/626127.

42.Hawkins D, Houeild N and Abrahamse H. Low level laser therapy (LLLT) as an effective therapeutic modality for delayed wound healing. *Annals of the New York Academy of Sciences*, 2005;1056(1):486-493.

43.Kana JS, Hutschenreiter G, Haina D and Waidelich W. Effect of low—power density laser radiation on healing of open skin wounds in rats. *Archives of Surgery*, 1981;116.3:293-296.

44.Anders JJ, Borke RC, Woolery SK and Van de

Merwe WP. Low power laser irradiation alters the rate of regeneration of the rat facial nerve. *Lasers in surgery and medicine*, 1993;13(1):72-82.

45. AlGhamdi KM, Kumar A and Moussa NA. Low-level laser therapy: a useful technique for enhancing the proliferation of various cultured cells. *Lasers in medical science*, 2012; 27(1):237-249.

46. Byrnes KR, Waynant RW, Ilev IK, Wu X, Barna L, Smith K, Heckert R, Gerst H and Anders JJ. Light

promotes regeneration and functional recovery and alters the immune response after spinal cord injury. *Lasers in surgery and medicine*, 2005;36(3):171-185.

47. Draper WE, Schubert TA, Clemmons RM and Miles A. Low-level laser therapy reduces time to ambulation in dogs after hemilaminectomy: a preliminary study. *Journal of Small Animal Practice*, 2012;53(8):465-469.

ارزیابی اثرات توأم پیوند موضعی سلول‌های بنیادی مزانشیمی خودی و درمان با لیزر کم‌توان در التیام جراحات حاد تجربی نخاع در رت

سید صدرا ایزدی^۱، محمدمهدی دهقان^{۱*}، داوود شریفی^۱، سید مهدی نصیری^۲، سیدحسین مرجانمهر^۴، مریم ذبیحی^۱، جواد اشرفی هلان^۵، حسام اکبرین^۶

^۱ گروه جراحی و رادیولوژی، دانشکده دامپزشکی، دانشگاه تهران، تهران، ایران

^۲ پژوهشکده تحقیقات زیست پزشکی، دانشگاه تهران، تهران، ایران

^۳ گروه بیماریهای داخلی، دانشکده دامپزشکی، دانشگاه تهران، تهران، ایران

^۴ گروه آسیب شناسی، دانشکده دامپزشکی، دانشگاه تهران، تهران، ایران

^۵ گروه پاتوبیولوژی، دانشکده دامپزشکی، دانشگاه تبریز، تبریز، ایران

^۶ گروه بهداشت و کنترل غذایی، دانشکده دامپزشکی، دانشگاه تبریز، تبریز، ایران

هدف- هدف از این مطالعه بررسی عملکردی و هیستوپاتولوژیک اثرات توأم پیوند خودی سلول‌های بنیادی مزانشیمی مغز استخوان و درمان با لیزر کم‌توان در ترمیم ضایعات حاد نخاع در رت بود.

طرح- مطالعه تجربی

حیوانات- تعداد ۲۸ سر رت نر بالغ نژاد ویستار.

روش کار: آسیب حاد نخاعی با ایجاد فشار بر روی نخاع بوسیله کاتتر بالون دار فوگارتی در محل مهره سینه ای ۸ یا ۹ القا شده و پس از ۷ روز در گروه اول، در محل ضایعه نخاعی سلول بنیادی مزانشیمی خودی تزریق گردید، در گروه دوم علاوه بر تزریق سلول بنیادی مزانشیمی خودی، تحت درمان با لیزر کم‌توان هم قرار گرفتند. در گروه سوم تنها درمان با لیزر کم‌توان را دریافت نمودند. در گروه چهارم هیچ درمانی انجام نشد (گروه کنترل). بررسی عملکردی رت‌ها در این گروه‌ها، در مدت ۵ هفته تا زمان بررسی هیستوپاتولوژیک بافت نخاع، انجام گردید.

نتایج- تغییرات آنالیز رفتاری در هفته‌های اول تا پنجم بین چهار گروه معنی دار بود ($P < 0.001$). اختلاف آماری معنی داری بین گروه سلول بنیادی مزانشیمی و گروه سلول بنیادی مزانشیمی بعلاوه لیزر کم‌توان؛ بین گروه سلول بنیادی مزانشیمی بعلاوه لیزر کم‌توان و گروه لیزر کم‌توان تنها؛ بین گروه سلول بنیادی مزانشیمی بعلاوه لیزر کم‌توان و گروه کنترل؛ بین گروه سلول بنیادی مزانشیمی و همچنین بین گروه لیزر کم‌توان تنها و گروه کنترل، در هفته‌های اول تا پنجم؛ مشاهده گردید. اختلاف آماری آنالیز رفتاری بین گروه سلول بنیادی مزانشیمی و لیزر کم‌توان تنها، فقط در هفته‌های اول و دوم معنی دار بود. نتایج هیستوپاتولوژیک نشان داد بهترین نتایج مربوط به گروه تحت درمان با سلول بنیادی مزانشیمی به همراه لیزر کم‌توان بود؛ بطوری که قسمت آسیب دیده نخاع تا حدی ترمیم و بازسازی شده بود. در ماده سفید، افزایش سلولهای گلیال، بازسازی سلولهای میلین و آکسونها مشاهده شد. ترمیم نورونها در شاخ پشتی مشاهده نشد و همانند گروه کنترل هیپوسولاریتی شدیدی در آنها دیده شد. در گروه درمان با سلول بنیادی، هنوز بقایای حضور ماکوفازها یا سلولهای میکروگلیال هضم کننده میلین دیده می‌شد. در گروه لیزر کم‌توان تنها هم تحریک سلولهای پشتیبان به تکثیر و هایپر تروفی و کمک به بازسازی نخاع مشهود بود، هرچند بازسازی کامل نبود.

نتیجه گیری و کاربرد بالینی- بررسی عملکردی و هیستوپاتولوژیک حاکی از نتایج بهتر در گروه درمان با سلول بنیادی مزانشیمی به همراه درمان با لیزر کم‌توان نسبت به سایر گروه‌ها بود، و می‌تواند نویدبخش یک درمان کارآمد برای آسیبهای نخاعی باشد؛ اگرچه مطالعات بیشتر و کاملتری در مقایسه با سایر روش‌های درمانی موجود ضروری به نظر می‌رسد.

کلمات کلیدی- آسیب نخاعی، سلول بنیادی مزانشیمی، درمان با لیزر کم‌توان، رت